Clayton County Public Schools
Allergy Action Plan (IHP)

Name: _____ **D.O.B.:** _____

Healthcare Provider: _____ **Provider's Phone:** _____

Parent/Guardian: _____ **Parent's Phone:** _____

Emergency Contact Name/Phone: _____ **School:** _____

Asthma: ☐ Yes (higher risk for a severe reaction) ☐ No

Extremely reactive to the following allergens: (please indicate) _____

THEREFORE:

☐ If checked, give epinephrine immediately for ANY symptoms if the allergen was *likely* eaten.

☐ If checked, give epinephrine immediately if the allergen was *definitely* eaten, even if no symptoms are noted.

Any SEVERE SYMPTOMS after suspected or known ingestion:

One or more of the following:

LUNG: Short of breath, wheeze, repetitive cough
HEART: Pale, blue, faint, weak pulse, dizzy, confused
THROAT: Tight, hoarse, trouble breathing/swallowing
MOUTH: Obstructive swelling (tongue and/or lips)
SKIN: Many hives over body

Or **combination** of symptoms from different body areas:

SKIN: Hives, itchy rashes, swelling (e.g., eyes, lips)
GUT: Vomiting, diarrhea, crampy pain

1. **INJECT EPINEPHRINE IMMEDIATELY**
2. Call 911
3. Begin monitoring (see box below)
4. Give additional medications:*
   -Antihistamine
   -Inhaler (bronchodilator) if asthma

*Antihistamines & inhalers/bronchodilators are not to be depended upon to treat a severe reaction (anaphylaxis). USE EPINEPHRINE.

MILD SYMPTOMS ONLY:

MOUTH: Itchy mouth
SKIN: A few hives around mouth/face, mild itch
GUT: Mild nausea/discomfort

1. **GIVE ANTIHISTAMINE**
2. Stay with student; alert healthcare professionals and parent
3. If symptoms progress (see above), USE EPINEPHRINE
4. Begin monitoring (see box below)

Medications/Doses

Epinephrine (brand and dose): _____

Antihistamine (brand and dose): _____

Other (e.g., inhaler-bronchodilator if asthmatic): _____

Monitoring

Stay with student; alert healthcare professionals and parent. Tell rescue squad epinephrine was given; request an ambulance with epinephrine. Note time when epinephrine was administered. A second dose of epinephrine can be given _____ minutes or more after the first if symptoms persist or recur. Treat student even if parents cannot be reached. See back/attached for auto-injection technique.

TURN FORM OVER **Form provided courtesy of the Food Allergy & Anaphylaxis Network (www.foodallergy.org) 9/2011**

SCHOOL MEDICATION CONSENT AND PROVIDER ORDER

_____ **This student is capable and approved to carry and self-administer the medicine(s) named on this form.**
_____ **This student is NOT approved to self-medicate.**
_____ **This student may be administered medication(s) by a school nurse or trained staff as directed.**

EPIPEN Auto-Injector and EPIPEN Jr Auto-Injector Directions

- **First, remove the EPIPEN Auto-Injector from the plastic carrying case**
- **Pull off the blue safety release cap**

- **Hold orange tip near outer thigh (always apply to thigh)**



- **Swing and firmly push orange tip against outer thigh. Hold on thigh for approximately 10 seconds. Remove the EPIPEN Auto-Injector and massage the area for 10 more seconds**

EPIPEN 2-PAK® | EPIPEN Jr 2-PAK®
(Epinephrine) Auto-Injectors 0.3/0.15mg

Adrenaclick™ 0.3 mg and Adrenaclick™ 0.15 mg Directions

Remove GREY caps labeled "1" and "2."

Place RED rounded tip against outer thigh, press down hard until needle penetrates. Hold for 10 seconds, then remove.

A food allergy response kit should contain at least two doses of epinephrine, other medications as noted by the student's physician, and a copy of this Food Allergy Action Plan.

A kit must accompany the student if he/she is off school grounds (i.e., field trip).

As the parent/guardian, I understand that the school, its employees and its agents shall incur no liability and shall be held harmless against any claims that may arise relating to the administration, supervision, training, or self-administration of medication.

Parent/Guardian Authorization	**Date**	**Physician/Healthcare Provider Authorization**	**Date**
Supervising Healthcare Professional	**Date**		

Form provided courtesy of the Food Allergy & Anaphylaxis Network (www.foodallergy.org) 9/2011